

CHICK STARTER CRUMBLES

As an aid in prevention of coccidiosis where immunity to coccidiosis is not desired.

ACTIVE DRUG INGREDIENT

Amprolium0.0125 %

Feeding Instructions

Feed as the sole ration to chicks from day-old until pullets are 6 weeks old. Always provide fresh, clean water.

WARNING

Withdraw 5 days before slaughter. Use as a sole source of amprolium.

Guaranteed Analysis

CRUDE PROTEIN (Min)	21.0%
CRUDE FAT (Min)	2.8%
CRUDE FIBER (Max)	3.0%
ASH (Max)	7.3%
PHOSPHORUS (Min)	0.8%
CALCIUM (Min)	1.1%
CALCIUM (Max)	1.3%
LYSINE (Min)	1.18%
SODIUM (Min)	0.2%
SODIUM (Max)	0.2%
SALT (Min)	0.3%
SALT (Max)	0.4%
SELENIUM (Min)	0.49ppm
SELENIUM (Max)	0.55ppm
METHIONINE (Min)	0.38%

List of Ingredients

Soybean Meal, Ground Corn, Wheat Ground, Wheat Millrun, Molasses, Calcium Carbonate, Dicalcium Phosphate, Blended Vegetable Oil with Ethoxyquin (a preservative), Amprolium 2.5%, Salt, Choline Chloride, Iron Sulfate, Manganous Oxide, Zinc Oxide, Ethoxyquin, Niacin, Vitamin E Supplement, Sodium Selenite, Copper Sulfate, D-Calcium Pantothenate, Ethylenediamine Dihydroiodide, Biotin, Menadione, Pyridoxine Hydrochloride, Riboflavin, Vitamin B12 Supplement, Folic Acid, Vitamin A Supplement, Vitamin D3 Supplement, Methionine Liquid.

50# K53555