

CHICK STARTER MASH

As an aid in prevention of coccidiosis where immunity to coccidiosis is not desired.

ACTIVE DRUG INGREDIENT

Amprolium0.0125%

Feeding Instructions

Feed as the sole ration to chicks from day-old until pullets are 6 weeks old. Always provide fresh, clean water.

WARNING

Withdraw 5 days before slaughter. Use as a sole source of Amprolium.

Guaranteed Analysis

CRUDE PROTEIN (Min)	21.0%
CRUDE FAT (Min)	3.1%
CRUDE FIBER (Max)	2.8%
ASH (Max)	7.0%
PHOSPHORUS (Min)	0.7%
CALCIUM (Min)	1.0%
CALCIUM (Max)	1.3%
LYSINE (Min)	1.20%
SODIUM (Min)	0.2%
SODIUM (Max)	0.2%
SALT (Min)	0.3%
SALT (Max)	0.4%
SELENIUM (Min)	0.43ppm
SELENIUM (Max)	0.48ppm
METHIONINE (Min)	0.37%

List of Ingredients

Ground Corn, Soybean Meal, Wheat Millrun, Calcium Carbonate, Dicalcium Phosphate, Molasses, Amprolium 2.5%, Blended Vegetable Oil with Ethoxyquin (a preservative), Salt, Choline Chloride, Iron Sulfate, Manganous Oxide, Zinc Oxide, Ethoxyquin, Niacin, Vitamin E Supplement, Sodium Selenite, Copper Sulfate, D-Calcium Pantothenate, Ethylenediamine Dihydroiodide, Biotin, Menadione, Pyridoxine Hydrochloride, Riboflavin, Vitamin B12 Supplement, Folic Acid, Vitamin A Supplement, Vitamin D3 Supplement, Sodium Sulfate, Sodium Sulfite, Potassium Sulfate, Sodium Bentonite, Propylene Glycol, Methionine Liquid.

25# K53602

50# K53605